



nephrogenic diabetes insipidus

Nephrogenic diabetes insipidus is a disorder of water balance. The body normally balances fluid intake with the excretion of fluid in urine. However, people with nephrogenic diabetes insipidus produce too much urine (polyuria), which causes them to be excessively thirsty (polydipsia). Affected individuals can quickly become dehydrated if they do not drink enough water, especially in hot weather or when they are sick.

Nephrogenic diabetes insipidus can be either acquired or hereditary. The acquired form is brought on by certain drugs and chronic diseases and can occur at any time during life. The hereditary form is caused by genetic mutations, and its signs and symptoms usually become apparent within the first few months of life.

Infants with hereditary nephrogenic diabetes insipidus may eat poorly and fail to gain weight and grow at the expected rate (failure to thrive). They may also be irritable and experience fevers, diarrhea, and vomiting. Recurrent episodes of dehydration can lead to slow growth and delayed development. If the condition is not well-managed, over time it can damage the bladder and kidneys leading to pain, infections, and kidney failure. With appropriate treatment, affected individuals usually have few complications and a normal lifespan.

Nephrogenic diabetes insipidus should not be confused with diabetes mellitus, which is much more common. Diabetes mellitus is characterized by high blood sugar levels resulting from a shortage of the hormone insulin or an insensitivity to this hormone. Although nephrogenic diabetes insipidus and diabetes mellitus have some features in common, they are separate disorders with different causes.

Frequency

The prevalence of nephrogenic diabetes insipidus is unknown, although the condition is thought to be rare. The acquired form occurs more frequently than the hereditary form.

Genetic Changes

The hereditary form of nephrogenic diabetes insipidus can be caused by mutations in at least two genes. About 90 percent of all cases of hereditary nephrogenic diabetes insipidus result from mutations in the *AVPR2* gene. Most of the remaining 10 percent of cases are caused by mutations in the *AQP2* gene. Both of these genes provide instructions for making proteins that help determine how much water is excreted in urine.

The acquired form of nephrogenic diabetes insipidus can result from chronic kidney disease, certain medications (such as lithium), low levels of potassium in the blood (hypokalemia), high levels of calcium in the blood (hypercalcemia), or an obstruction of the urinary tract.

The kidneys filter the blood to remove waste and excess fluid, which are stored in the bladder as urine. The balance between fluid intake and urine excretion is controlled by a hormone called vasopressin or antidiuretic hormone (ADH). ADH directs the kidneys to concentrate urine by reabsorbing some of the water into the bloodstream. Normally, when a person's fluid intake is low or when a lot of fluid is lost (for example, through sweating), increased levels of ADH in the blood tell the kidneys to make less urine. When fluid intake is adequate, lower levels of ADH tell the kidneys to make more urine.

Mutations in the *AVPR2* or *AQP2* genes prevent the kidneys from responding to signals from ADH. Chronic kidney disease, certain drugs, and other factors can also impair the kidneys' ability to respond to this hormone. As a result, the kidneys do not reabsorb water as they should, and the body makes excessive amounts of urine. These problems with water balance are characteristic of nephrogenic diabetes insipidus.

Inheritance Pattern

When nephrogenic diabetes insipidus results from mutations in the *AVPR2* gene, the condition has an X-linked recessive pattern of inheritance. The *AVPR2* gene is located on the X chromosome, which is one of the two sex chromosomes. In males (who have only one X chromosome), one altered copy of the gene in each cell is sufficient to cause the condition. In females (who have two X chromosomes), a mutation usually has to occur in both copies of the gene to cause the disorder. However, some females who carry a single mutated copy of the *AVPR2* gene have features of nephrogenic diabetes insipidus, including polyuria and polydipsia. A characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons.

When nephrogenic diabetes insipidus is caused by mutations in the *AQP2* gene, it can have either an autosomal recessive or, less commonly, an autosomal dominant pattern of inheritance. In autosomal recessive inheritance, both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition. In autosomal dominant inheritance, one mutated copy of the *AQP2* gene in each cell is sufficient to cause the disorder.

Other Names for This Condition

- ADH-resistant diabetes insipidus
- congenital nephrogenic diabetes insipidus
- diabetes insipidus renalis
- diabetes insipidus, nephrogenic

- NDI
- vasopressin-resistant diabetes insipidus

Diagnosis & Management

These resources address the diagnosis or management of nephrogenic diabetes insipidus:

- GeneReview: Nephrogenic Diabetes Insipidus
<https://www.ncbi.nlm.nih.gov/books/NBK1177>
- Genetic Testing Registry: Nephrogenic diabetes insipidus
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0162283/>
- Genetic Testing Registry: Nephrogenic diabetes insipidus, X-linked
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1563705/>
- Genetic Testing Registry: Nephrogenic diabetes insipidus, autosomal
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1563706/>
- MedlinePlus Encyclopedia: ADH
<https://medlineplus.gov/ency/article/003702.htm>
- MedlinePlus Encyclopedia: Diabetes Insipidus - Nephrogenic
<https://medlineplus.gov/ency/article/000511.htm>

These resources from MedlinePlus offer information about the diagnosis and management of various health conditions:

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: ADH
<https://medlineplus.gov/ency/article/003702.htm>
- Encyclopedia: Diabetes Insipidus
<https://medlineplus.gov/ency/article/000377.htm>
- Encyclopedia: Diabetes Insipidus - Nephrogenic
<https://medlineplus.gov/ency/article/000511.htm>
- Health Topic: Diabetes Insipidus
<https://medlineplus.gov/diabetesinsipidus.html>

Genetic and Rare Diseases Information Center

- Nephrogenic diabetes insipidus
<https://rarediseases.info.nih.gov/diseases/7178/nephrogenic-diabetes-insipidus>

Additional NIH Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: Diabetes Insipidus
<https://www.niddk.nih.gov/health-information/kidney-disease/diabetes-insipidus>
- National Institute of Diabetes and Digestive and Kidney Diseases: The Kidneys and How They Work
<https://www.niddk.nih.gov/health-information/kidney-disease/kidneys-how-they-work>

Educational Resources

- Boston Children's Hospital
<http://www.childrenshospital.org/conditions-and-treatments/conditions/diabetes-insipidus>
- Disease InfoSearch: Nephrogenic diabetes insipidus
<http://www.diseaseinfosearch.org/Nephrogenic+diabetes+insipidus/5139>
- Disease InfoSearch: Nephrogenic diabetes insipidus, autosomal
<http://www.diseaseinfosearch.org/Nephrogenic+diabetes+insipidus%2C+autosomal/8974>
- MalaCards: diabetes insipidus, nephrogenic
http://www.malacards.org/card/diabetes_insipidus_nephrogenic

- Merck Manual Consumer Version
<http://www.merckmanuals.com/home/kidney-and-urinary-tract-disorders/disorders-of-kidney-tubules/nephrogenic-diabetes-insipidus>
- Orphanet: Nephrogenic diabetes insipidus
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=223

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD)
<https://rarediseases.org/rare-diseases/nephrogenic-diabetes-insipidus/>
- Resource list from the University of Kansas Medical Center
<http://www.kumc.edu/gec/support/diabetes.html>

GeneReviews

- Nephrogenic Diabetes Insipidus
<https://www.ncbi.nlm.nih.gov/books/NBK1177>

Genetic Testing Registry

- Nephrogenic diabetes insipidus
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0162283/>
- Nephrogenic diabetes insipidus, autosomal
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1563706/>
- Nephrogenic diabetes insipidus, X-linked
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1563705/>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22nephrogenic+diabetes+insipidus%22>

Scientific articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Diabetes+Insipidus,+Nephrogenic%5BMAJR%5D%29+AND+%28nephrogenic+diabetes+insipidus%5BTIAB%5D%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- DIABETES INSIPIDUS, NEPHROGENIC, AUTOSOMAL
<http://omim.org/entry/125800>
- DIABETES INSIPIDUS, NEPHROGENIC, X-LINKED
<http://omim.org/entry/304800>

Sources for This Summary

- Bichet DG. Nephrogenic diabetes insipidus. *Adv Chronic Kidney Dis.* 2006 Apr;13(2):96-104. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16580609>
 - Fujiwara TM, Bichet DG. Molecular biology of hereditary diabetes insipidus. *J Am Soc Nephrol.* 2005 Oct;16(10):2836-46. Epub 2005 Aug 10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16093448>
 - GeneReview: Nephrogenic Diabetes Insipidus
<https://www.ncbi.nlm.nih.gov/books/NBK1177>
 - Khanna A. Acquired nephrogenic diabetes insipidus. *Semin Nephrol.* 2006 May;26(3):244-8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16713497>
 - Knoers NV, Deen PM. Molecular and cellular defects in nephrogenic diabetes insipidus. *Pediatr Nephrol.* 2001 Dec;16(12):1146-52. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11793119>
 - Morello JP, Bichet DG. Nephrogenic diabetes insipidus. *Annu Rev Physiol.* 2001;63:607-30. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11181969>
 - van Lieburg AF, Knoers NV, Monnens LA. Clinical presentation and follow-up of 30 patients with congenital nephrogenic diabetes insipidus. *J Am Soc Nephrol.* 1999 Sep;10(9):1958-64.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10477148>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/nephrogenic-diabetes-insipidus>

Reviewed: April 2010

Published: January 24, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services